



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 18, 2015

Shannon Yanocha  
Product Registration Manager  
FMC Corporation  
1735 Market Street  
Philadelphia, PA 19103

Subject: Label Amendment – Update the label  
Product Name: F7127 SE HERBICIDE  
EPA Registration Number: 279-3337  
Application Date: 1/20/15  
Decision Number: 499487

Dear Ms Yanocha:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 279-3337  
Decision No. 499487

with FIFRA section 6. If you have any questions, you may contact Dominic Schuler at 703-347-0260 or via email at [schuler.dominic@epa.gov](mailto:schuler.dominic@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending from the end of the name.


Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

# F7127 SE HERBICIDE

EPA Reg. No. 279-3337

EPA Est. No. XXX



Active Ingredients:	By Wt.
Carfentrazone-ethyl* .....	3.53%
Sulfentrazone** .....	31.77%
Other Ingredients: .....	64.70%
<b>Total:</b> .....	<b>100.00%</b>

\*F7127 SE HERBICIDE contains 0.35 pounds per US gallon of the active ingredient Carfentrazone-ethyl.

\*\*F7127 SE HERBICIDE contains 3.15 pounds per US gallon of the active ingredient Sulfentrazone.

U.S. Patent Pending

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

### FIRST AID

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

F7127 SE HERBICIDE IS FORMULATED AND PACKAGED IN USA.

FROM 

1735 Market Street  
Philadelphia, PA 19103

<b>TABLE OF CONTENTS</b>	
<b>Section Title</b>	<b>Page</b>
Active Ingredient	1
First Aid Statements	1
Hotline Number	1
Table of Contents	2
Precautionary Statements	2
Personal Protective Equipment (PPE)	2
User Safety Recommendations	3
Environmental Hazards	3
Physical/Chemical Hazards	3
Conditions of Sale and Limitation of Warranty and Liability	3
Resistance Management	3
Directions for Use	4
Agricultural Use Requirements	4
Restricted Entry Interval (REI)	4
Storage and Disposal	4
Product Information	5
Product Application Instructions	5
Environmental and Soil Factors Influencing F7127 SE Herbicide Applications	6
Soil Classification Chart	6
Mixing and Loading Instructions	6
Application Information	7
Spray Drift Reduction Advisory	8
Spray Equipment Clean-out	9
Maximum Allowable F7127 SE Herbicide Use Per Acre Per 12 Month Period	9
Rate Conversion Chart	10
Crop Rotation Intervals	10
Replanting Instructions	11
Postemergent Weeds Controlled	11
Cabbage (Transplanted only)	13
Corn (Field Corn, Seed Corn, Popcorn)	14
Dry Shelled Beans and Peas	15
Fallow Systems	17
Flax	18
Horseradish	19
Lima Beans, Succulent (Tennessee only)	20
Peanuts	21
Potatoes	22
Soybeans	24
Sugarcane	25
Sunflowers	26
Tobacco (Burley, Flue-Cured and Dark)	27
Sod Production	29
Permanent Crops	31
Apples	31
Citrus Fruit	31
Tree Nuts	31
Grapes	31
Berries	31
Weed Control Information	32
Label Tracking Information	35

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

#### **Caution**

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

### **Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear: protective eyewear (goggles or face shield), long-sleeved shirt and long pants, chemical resistant gloves made of waterproof material such as polyethylene or polyvinyl chloride, and shoes plus socks.**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## **User Safety Recommendations**

### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **Environmental Hazards**

This pesticide is toxic to algae, marine/estuarine invertebrates, and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

### **Groundwater Advisory**

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Do not use on coarse soils classified as sand which have less than 1% organic matter.**

### **Surface Water Advisory**

This product can contaminate surface water through spray drift. Under some conditions, this product may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

## **Physical/Chemical Hazards**

Combustible. Do not use or store near heat or open flame.

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product.

If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and, to the extent consistent with applicable law, buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

## **RESISTANCE MANAGEMENT**

Some weeds are known to develop resistance to herbicides that have been used repeatedly. While the development of resistance is well understood, it is not easily predicted. Use herbicides in conjunction with resistance management strategies in the area. Consult the local or State agricultural advisors for details. If weed resistance should develop in the area, this product used alone may not continue to provide sufficient levels of weed control. If the reduced levels of control cannot be attributed to improper application timing, unfavorable weather conditions or abnormally high weed pressure, a resistant strain may have developed.

To reduce the potential for weed resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the listed rates and in accordance with the use directions. Do not use less than listed label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. For optimum performance, scout fields carefully and begin applications when weeds are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants, chemical resistant gloves, and shoes plus socks.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

#### Pesticide storage

Store product in original container only, away from other pesticides, fertilizer, food or feed. Store in a cool dry place and avoid excess heat. Do not store below 32F degrees.

#### In Case of Spill

Avoid contact. Isolate areas and keep out animals and unprotected persons.

#### To Confine Spills

Dike surrounding area, sweep up spillage, Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a large holding container. Identify contents per required hazardous waste labeling regulations.

#### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

#### Container Handling

**Nonrefillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: (For containers greater than 5 gallons) Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. (For containers 5 gallons or less) Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Triple rinse (or equivalent). Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Containers** - Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## PRODUCT INFORMATION

F7127 SE HERBICIDE is a selective herbicide that provides postemergent contact and soil residual weed control. F7127 SE HERBICIDE may be applied as a burndown prior to planting, early preplant, or as a preemergent application before or after weed emergence for control of susceptible broadleaf weeds. F7127 SE HERBICIDE is a 3.5 pound per gallon suspoemulsion containing the active ingredients carfentrazone-ethyl and sulfentrazone. Applications of F7127 SE HERBICIDE must be made before crop seed germination to prevent injury to the emerging crop seedlings. When applications after planting are delayed, injury may occur if seeds are germinating or if they are located near the soil surface.

Observe the most restrictive of all instructions, crop restrictions, mixing directions, application precautions, replanting directions, rotational crop guidelines and other label information of each product when tank mixing with F7127 SE HERBICIDE. In addition to application information, refer to the specific directions of use for a particular crop/use pattern as set forth below.

### Proper Handling Instructions

This product must not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

**Do not apply this product through any type of irrigation system.**

**Do not use flood irrigation to apply or incorporate this product.**

This product must be used in a manner which will prevent back siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

## PRODUCT APPLICATION INSTRUCTIONS

F7127 SE HERBICIDE is to be mixed with water, liquid fertilizer, or mixtures of water and liquid fertilizer and applied in fallow systems or as a preplant burndown and/or preemergence herbicide to labeled crops. F7127 SE HERBICIDE provides postemergent contact and soil residual control of susceptible weed species.

Emerged, susceptible broadleaf weeds are easiest to control when they are small (less than 3 inches tall) and actively growing. Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved postemergent weed control will be poor. Always use the higher application rate of this product, for the appropriate soil texture and organic matter, when weed growth is dense or heavy, or when weeds are growing in an undisturbed or non-cultivated area. Reduced weed control may occur if weeds are experiencing drought stress, disease or insect damage, or when weeds are thickly covered with dust. For control of weeds not listed on this label F7127 SE HERBICIDE may be tank-mixed with other herbicides such as glyphosate, paraquat and glufosinate. Read and follow all manufacturers' label directions for the companion herbicide(s) and follow the most restrictive instructions for use. The use of a quality spray adjuvant is required for optimum control of emerged weeds. Refer to the individual crop sections of this label for specific adjuvant type and use rates.

The residual activity of F7127 SE HERBICIDE applications requires adequate moisture for herbicidal activation. The amount of residual activity is dependent on several factors. These factors include, but are not limited to, existing soil moisture at application, soil type, organic matter, and tillth. Where irrigation is not available and rainfall has not provided activation, particularly for surface applications of F7127 SE HERBICIDE, a shallow incorporation (less than 2") is recommended for destruction of any existing weeds and to incorporate F7127 SE HERBICIDE. Herbicide incorporation will initiate the process of activation with existing soil moisture. In circumstances where rainfall has not occurred and/or irrigation is not possible, alternative or additional weed management practices may be required.

Under normal growing conditions, F7127 SE HERBICIDE exhibits excellent crop safety. Soil applications of F7127 SE HERBICIDE must be made before crop seed germination to prevent injury to the emerging crop seedlings. F7127 SE HERBICIDE applied after crop emergence will cause severe injury to the crop. Poor growing conditions, such as excessive moisture, cool temperatures, and soil compaction or the presence of various pathogens may impact seedling or plant vigor. Under these conditions, the active ingredients in F7127 SE HERBICIDE can contribute to crop response. Refer to the specific directions of use for a particular crop/use pattern as set forth below for additional information.

## ENVIRONMENTAL AND SOIL FACTORS INFLUENCING F7127 SE HERBICIDE APPLICATIONS

Do not apply to soils classified as sand with less than 1% organic matter.

The user is required to read and follow the specific F7127 SE HERBICIDE use directions and restrictions for each crop as defined in subsequent sections of this label. The user is cautioned that some crops and weeds respond differently to F7127 SE HERBICIDE. This response is governed by the F7127 SE HERBICIDE application rate, various soil factors and inherent crop sensitivity. See individual crop use sections for specific directions on the use of F7127 SE HERBICIDE for optimum weed control and crop safety results in each crop.

### INFLUENCE OF CLAY, SOIL TYPE, AND PH ON F7127 SE HERBICIDE USE RATES AND CROP RESPONSE

Following an application of F7127 SE HERBICIDE to soil, germinating seeds and seedlings take up F7127 SE HERBICIDE from the soil solution. The amount of F7127 SE HERBICIDE in the soil solution, and available for weed uptake, is determined primarily by soil type, organic matter, and soil pH. F7127 SE HERBICIDE adsorbs to the clay and organic matter fractions of soils; effectively limiting the amount of active ingredient immediately available to control weeds. Soils typically increase in clay content through the series from coarse to fine as noted in the following Soil Classification Chart (Table 1).

**Table 1. SOIL CLASSIFICATION CHART**

COARSE	MEDIUM	FINE
Sand	Sandy clay loam	Silty clay loam
Loamy sand	Sandy clay Loam	Silty clay
Sandy loam	Silt loam	Clay loam
	Silt	Clay

Soil organic matter content can vary widely and independently of soil type and requires an accurate analysis of representative soil samples to determine its content.

### Do not use this product on coarse soils classified as sand which have less than 1% organic matter.

Soil pH also exerts a dramatic affect on F7127 SE HERBICIDE availability in the soil solution. As soil pH increases, F7127 SE HERBICIDE availability increases. Accurate soil pH information will require an accurate analysis of representative soil samples.

The total amount of F7127 SE HERBICIDE available, in any given soil, is determined by the interaction of soil type (clay content), % organic matter, and pH. The application timing (relative to the emergence of the crop and weeds) and amount of rainfall and/or irrigation received will ultimately determine, in conjunction with the soil parameters and pH, the amount of F7127 SE HERBICIDE in soil solution.

Irrigation with highly alkaline water (high pH) following a F7127 SE HERBICIDE soil application can also significantly increase the amount of F7127 SE HERBICIDE available in the soil solution. Irrigation with water having a pH greater than 7.5 could result in adverse crop response. This response will ultimately depend on initial F7127 SE HERBICIDE application rate, timing, amount and pH of irrigation water and sensitivity of the crop and its growth stage when irrigated. The risk of adverse crop response will lessen with the advance in growth stage among most crops.

The following Crop Specific Use Directions have been designed with specific F7127 SE HERBICIDE instructions for each crop based on the soil type, soil organic matter, and soil pH interactions described above. The user is cautioned that crop tolerance and weed control performance are based on strict adherence to these use directions.

## MIXING AND LOADING INSTRUCTIONS

Water or liquid fertilizer solutions may be used as the carrier for F7127 SE HERBICIDE when applied alone or in tank mixtures with other registered herbicides. Conduct a jar test to determine the compatibility of F7127 SE HERBICIDE and the fertilizer solution. When mixing with fertilizer solutions it is important to premix F7127 SE HERBICIDE in clear water. See directions for applying F7127 SE HERBICIDE alone with liquid fertilizer in Application Information.

A crop oil concentrate, methylated seed oil, nonionic surfactant (NIS) wetting agent labeled, or other equivalent adjuvant labeled for use with herbicides is required for optimum control of emerged weeds. Read and follow all applicable use directions, precautions and restrictions on the surfactant label.

### F7127 SE HERBICIDE Applied Alone

Select the proper F7127 SE HERBICIDE application rate from the following tables in the crop section of this label. Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of F7127 SE HERBICIDE for acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Apply the F7127 SE HERBICIDE spray mixture immediately after mixing.

Do not store spray mixture.

Do not prepare spray mixtures in nurse tanks.

### **F7127 SE HERBICIDE Applied in Tank Mix Combination**

Select the proper F7127 SE HERBICIDE application rate from TIMING AND METHOD OF APPLICATION section of label. Read and follow the most restrictive of all applicable use directions, precautions and restrictions on the respective tank mix product labels. To ensure product compatibility, a jar test should be conducted before large volume mixing (see MIXTURE COMPATIBILITY TESTING chart below). Provided the jar test indicates the mixture is compatible, prepare the tank mixture as follows.

Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of F7127 SE HERBICIDE for the acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, F7127 SE HERBICIDE and other liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Allow time for complete mixing and dispersion after each addition, adding water as necessary. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Use F7127 SE HERBICIDE tank mixtures immediately after mixing.

Do not store tank mixtures.

Do not prepare spray mixtures in nurse tanks.

### **F7127 SE HERBICIDE Applied Alone with Liquid Fertilizer**

When adding F7127 SE HERBICIDE to a liquid fertilizer carrier, premix F7127 SE HERBICIDE in clear water before adding to fertilizer solution. Adding F7127 SE HERBICIDE to fertilizer mixtures without first mixing with water can result in incompatibility.

Fill the spray tank one-half full with fertilizer solution. With agitator operating, add the F7127 SE HERBICIDE slurry to the spray tank. Use a minimum of one gallon of water for each container of F7127 SE HERBICIDE. Then add slurry to the spray tank through a 20-35 mesh screen. Rinse container used for pre-mixing and add rinsate to the spray tank. Complete filling the sprayer tank with fertilizer. Maintain agitation during filling, mixing and application. Use F7127 SE HERBICIDE spray mixture immediately after mixing. Do not store mixture.

Do not prepare spray mixtures in nurse tanks.

### **Jar Testing Fertilizer Spray Mixtures**

Applications of F7127 SE HERBICIDE alone, or with recommended tank mixtures, in conjunction with clear liquid fertilizer solutions (28-32% nitrogen only) may be used unless use directions specifically state otherwise. Test small quantities for compatibility by the following procedure before mixing in full spray tank quantities.

- 1) Add 1 pint of fertilizer solution in a quart jar.
- 2) Add the appropriate amount of herbicide based on the MIXTURE COMPATIBILITY table below. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).
- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again after 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, and then follow directions noted below.

### **MIXTURE COMPATIBILITY TESTING**

<b>Herbicide Type</b>	<b>Herbicide Field Use Rate</b>	<b>Amount Herbicide Added Per Pint</b>
Wettable Powder or Dry Flowable	0.5 pound	0.75 teaspoon
	1.0 pound	1.50 teaspoons
	2.0 pounds	3.00 teaspoons
	3.0 pounds	4.50 teaspoons
Emulsified Concentrates	1.0 pint	0.5 teaspoon
Liquid Flowables	1.0 quart	1.0 teaspoon
	2.0 quarts	2.0 teaspoons
	3.0 quarts	3.0 teaspoons

\*Based on a spray volume of 25 gallons per acre. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

### **Adjuvant Requirements**

The use of methylated seed oil (MSO) or a crop oil concentrate (COC) adjuvant, labeled for use with herbicides, is required for optimum control of emerged weeds. A nonionic surfactant adjuvant and water conditioning agent is recommended when F7127 SE HERBICIDE is tank-mixed with glyphosate. Read and follow all applicable use directions, precautions and restrictions on the surfactant label.

## **APPLICATION INFORMATION**

### **Ground Application**

Use a boom and/or nozzle sprayer equipped with the appropriate nozzles and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a

minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population. Thorough coverage is essential for control of susceptible broadleaf weeds. Be aware that overlaps and slower ground speeds while starting, stopping, or turning while spraying may result in excessive application and subsequent crop response.

Continuous agitation is required until all spray mixture has been applied. Avoid swath overlaps. Shut off spray booms while turning, slowing or stopping, as over application may result. Do not allow F7127 SE HERBICIDE spray mixtures to sit overnight as settling of product and difficulty of re-suspending may occur.

To avoid injury to sensitive crops, spray equipment used for F7127 SE HERBICIDE applications must be drained and thoroughly cleaned with water plus ammonia before being used to apply other products. See Spray Clean-out section.

**Avoid all direct, and/or indirect spray contact with non-target plants.** Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants to minimize exposure. Do not apply when wind speed favors drift beyond the area of treatment.

#### **Aerial Application**

F7127 SE HERBICIDE may be applied by air using properly calibrated nozzle types and arrangements that will provide optimum coverage while producing minimal amounts of fine droplets. Apply sufficient spray volume to achieve adequate coverage. Apply a minimum of five (5) gallons of finished spray per acre. Do not apply when wind speed favors drift beyond the area intended for treatment.

#### **Runoff and Wind Erosion Precautions**

**Do not apply under conditions which favor runoff or wind erosion of soil containing F7127 SE HERBICIDE to non-target areas.**

**To prevent off-site movement due to runoff or wind erosion:**

- Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, allow the soil surface to be settled by rainfall or irrigation.
- Do not apply to impervious substrates such as paved or highly compacted surfaces or frozen or snow covered ground.
- Do not apply to soils when saturated with water.
- Do not use tail water from the first flood or furrow irrigation of treated fields to treat non target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

## **SPRAY DRIFT REDUCTION ADVISORY**

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops.

Where States and local governments have more stringent regulations, they must be observed.

#### **Droplet Size Information**

Reduce drift potential by applying large droplets. The optimum drift management strategy is to apply the largest droplets that will provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift when applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity and Temperature Inversions).

**VMD** – VMD is the expression of the droplet size of the spray cloud. The VMD value means that 50% of the droplets are larger than the expressed value and 50% of the droplets are smaller than the expressed value. Optimum spray clouds should be 450 microns with fewer than 10% of the droplets being 200 microns or smaller.

**Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows usually produce larger droplets.

**Pressure** – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types narrower spray angles produce larger droplets. Consider using low drift nozzles.

**Application Height** – Making applications at the lowest height practical reduces exposure of spray droplets to evaporation and wind movement.

**Swath Adjustment** – Swath adjustment distance must increase with increasing drift potential (higher wind, smaller droplets, etc.)

**Wind** – Drift potentials are lowest between wind speeds of 3 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Applications in wind conditions outside of this range could increase the risk of off-target effects and should be avoided. Note that local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity** - When making applications in conditions of low relative humidity set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions** – Do not apply F7127 SE HERBICIDE during temperature inversions because the drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the following morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or a smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicate an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas** – Only apply when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitats for threatened or endangered species and non-target crops).

**Off-Target Movement of F7127 SE HERBICIDE** – Drift of dilute spray mixtures containing F7127 SE HERBICIDE must be prevented. Observation of the preceding environmental conditions, correct application equipment design, calibration and application practices will significantly diminish the risk of off-target spray drift. F7127 SE HERBICIDE can cause significant symptomology by drift on to sensitive crops and other plants. This symptomology may manifest initially as discreet, localized spots where contact by F7127 SE HERBICIDE drift mixtures. Depending on concentration of the spray solution and droplet size (effectively determining the concentration of F7127 SE HERBICIDE) and also depending on the inherent sensitivity of the plants involved, these spots or lesions may not coalesce. These effects will usually not have lasting effects on plant growth, but will likely reduce the value of affected fruit of foliage where grade or quality is associated with appearance. In severe drift instances with particularly sensitive crops, defoliation of affected foliage could result. Failure to follow these guidelines and environmental prohibitions that then result in off-target movement or drift of F7127 SE HERBICIDE on to unintended crops or plants, irrespective of severity, constitutes misapplication of this product. FMC accepts no responsibility or liability for potential crop effects that may result from such misapplication of F7127 SE HERBICIDE.

## SPRAY EQUIPMENT CLEAN-OUT

**After spraying F7127 SE HERBICIDE and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure.**

1. Drain sprayer tank, hoses, and spray boom and thoroughly rinse the inside of the sprayer tank with clean water to remove sediment and residues. Thoroughly flush sprayer hoses, boom and nozzles with clean water.
2. Fill the tank 1/2 full with clean water, and add appropriate detergent or ammonia (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom, and nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the cleaning solution is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Before using the sprayer, drain the spray system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately with the detergent or ammonia solution.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State and local regulations and guidelines.

Do not drain or flush equipment on or near desirable trees or plants.

Do not contaminate any body of water including irrigation water that may be used on other crops.

Should small quantities of F7127 SE HERBICIDE remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. FMC accepts no liability for any effects due to inadequately cleaned equipment.

## MAXIMUM ALLOWABLE F7127 SE HERBICIDE USE PER ACRE PER 12 MONTH PERIOD\*

Refer to the crop section of this label for specific product use directions.

Crop	Fluid Ounces F7127 SE HERBICIDE Per Acre	Pounds Active F7127 SE HERBICIDE** Per Acre
Apples	15.2	0.41
Berries (Crop Group 13-07)	15.2	0.41
Cabbage	15.2	0.41
Citrus (Crop Group 10)	15.2	0.41
Corn	10.2	0.28

Dry peas & beans	10.2	0.28
Fallow	10.2	0.28
Flax	10.2	0.28
Grapes	15.2	0.41
Horseradish	10.2	0.28
Lima beans, succulent (Tennessee only)	7.6	0.21
Soybeans	8.5	0.23
Sunflowers	10.2	0.28
Peanut	12.2	0.33
Potato	10.2	0.28
Sod production	15.2	0.41
Sugarcane	15.2	0.41
Tobacco	15.2	0.41
Tree Nuts (Crop Group 14)	15.2	0.41

\*The total allowed usage per twelve-month period includes all applications made to the field per twelve-month interval. This includes fallow treatments, burndown treatments, planting time and all in-season treatments. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

\*\* Based on total active carfentrazone-ethyl and sulfentrazone

## RATE CONVERSION CHART

F7127 HERBICIDE		SE	CARFENTRAZONE-ETHYL		SULFENTRAZONE	
Product fl oz/A	lb ai*		Product fl oz/A**	lb ai	Product fl oz/A***	lb ai
3.75	0.10		0.65	0.01	2.9	0.09
5.75	0.15		1.0	0.015	4.5	0.14
8.50	0.23		1.5	0.02	6.7	0.21
10.20	0.28		1.8	0.03	8.0	0.25
15.25	0.41		2.7	0.04	12.0	0.37

\* Total pounds active of sulfentrazone + carfentrazone-ethyl

\*\* Based on Aim 2EC formulation

\*\*\* Based on SPARTAN 4F formulation

## CROP ROTATIONAL INTERVALS

Shown below are the minimum intervals in months from the time of F7127 SE HERBICIDE application until F7127 SE HERBICIDE treated soil may be replanted with the crops listed. When F7127 SE HERBICIDE is tank mixed with other herbicide(s), refer to all those labels for re-cropping instructions, following the intervals that are the most restrictive. For crops not listed, the interval is 12 months in addition to a successful field bioassay.

The field bioassay is a test strip of the intended crop planted across the previously treated field and grown to maturity. The test strip should include low spots, knolls, and variable pH and soil types. If crop responses are not observed, the crop may be planted the following year.

### CROP ROTATION INTERVALS\*

CROP	INTERVAL (Months)
Apples	Anytime
Alfalfa	12
Barley	4
Berries (Crop subgroup 13-07)	Anytime
Cabbage (transplant only)	Anytime
Canola, Crambe	24
Citrus (Crop Group 10)	Anytime
Corn, field	Anytime
Corn, pop	Anytime
Corn, seed	6
Corn, sweet	12
Cotton	18 or 12***
Dry Shell Peas & Beans	Anytime
Flax	Anytime
Grapes	Anytime
Horseradish	Anytime

Lima Beans (Tennessee only)	Anytime
Peanuts	Anytime
Peppermint	Anytime
Potatoes	Anytime
Rice	10
Rye	4
Sorghum	10**
Soybeans	Anytime
Spearmint	Anytime
Sugar Beets	36 or 24†
Sugarcane	Anytime
Sunflowers	Anytime
Sweet Potatoes	12
Tobacco	Anytime
Tomato (Transplanted)	Anytime
Tree Nuts (Crop group 14)	Anytime
Triticale	4
Turf	Anytime
Wheat	4

\* For all other crops not listed, the rotation interval is a minimum of 12 months with a successful bioassay.

\*\* 18 month rotation for rates above 10.2 fluid ounces per acre. Crops that have rotational intervals greater than 12 months after a F7127 SE HERBICIDE application are the result of crop injury concerns. Only plant after a successful bioassay.

\*\*\* Cotton may be planted after 12 months where F7127 SE HERBICIDE was applied at rates 8 oz/acre or less and meets the following conditions:

- Medium and fine soils
- pH <7.2
- Rainfall or irrigation must exceed 15" after application of F7127 SE HERBICIDE to rotate to cotton

†Sugar beets can be planted after 24 months with a successful bioassay.

## REPLANTING INSTRUCTIONS

If the initial planting of labeled crops fails to produce a uniform stand, only labeled crops for F7127 SE HERBICIDE or the tank mix partner; whichever is most restrictive, may be replanted. Do not retreat fields with a second application of F7127 SE HERBICIDE or other herbicide containing sulfentrazone. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals that are inconsistent with the CROP ROTATION INTERVALS on this label. When replanting use minimum soil tillage to preserve the herbicide barrier and achieve maximum weed control.

## POSTEMERGENT WEEDS CONTROLLED

**Pre-Plant Burndown** (Refer to individual crop sections for preemergent weeds controlled).

When used as directed, F7127 SE HERBICIDE will provide postemergent control of the following weeds (less than 3 inches tall) in a conventional till program. For complete no-till postemergence burndown control of the weeds listed and other no-till weeds a tank-mix of F7127 SE HERBICIDE and glyphosate or other labeled burndown herbicide is recommended.

Weeds Controlled	F7127 SE HERBICIDE use rate fluid ounce (lb ai) per acre
Lambsquarters (up to 3 inches tall)	3.75 (0.10)
Morningglory, Ivyleaf (up to 3 leaves)	
Morningglory, pitted (up to 3 leaves)	
Nightshade, Eastern black	
Pigweed, redroot	
Velvetleaf	
Waterhemp (up to 2 inches tall)	
Weeds Controlled	F7127 SE HERBICIDE use rate fluid ounce (lb ai) per acre
All the weeds controlled at 3.75 fluid ounces per acre (0.10 lb/acre) plus the weeds listed below:	4.75 (0.13)
Cheeseweed	
Filaree, redstem	
Flixweed	
Lambsquarters, common	

Mallow, common	
Morningglory, entireleaf	
Morningglory, ivyleaf	
Morningglory, pitted	
Morningglory, scarlet	
Nightshade, hairy	
Pennycress, field	
Pigweed, smooth	
Sesbania, hemp	
Smartweed (PA), seedling	
Tansymustard	
Waterhemp	
<b>Weeds Controlled</b>	<b>F7127 SE HERBICIDE use rate fluid ounce (lb ai) per acre</b>
All the weeds controlled at 4.75 fluid ounces per acre (0.13 lb/acre) plus the weeds listed below:	
Amaranth, spiny	
Anoda, spurred	
Bedstraw, catchweed	
Buffalobur	
Carpetweed	
Cocklebur	
Copperleaf, hophornbeam	
Cotton, GMO varieties	
Cotton, volunteer	
Dayflower	
Eclipta	
Fiddleneck, coast	
Groundcherry, smooth (seedling)	
Groundcherry, Wright's	
Jimsonweed	
Kochia	
Rocket, London	
Morningglory, ivyleaf	
Morningglory, tall	
Nightshade, American black	
Nightshade, black	
Shepherdspurse	
Spiderwort, tropical	
Thistle, Russian	
Wallflower, bushy	
<b>Weeds Controlled</b>	<b>F7127 SE HERBICIDE use rate fluid ounce (lb ai) per acre</b>
All the weeds controlled at 6.0 fluid ounces per acre (0.16 lb/acre) plus the weeds listed below:	
Amaranth, Palmer (up to 4 inches tall)	
Ammania, purple	
Buckwheat, wild	
Burclover	
Filaree, broadleaf	
Filaree, white	
Lettuce, prickly	
Mallow, Venice (up to 2 inches tall)	

6.0 (0.16)

8.5 (0.23) – 15.2 (0.41)

Meadowfoam	
Mustard spp.	
Redmaids	
Spurry, corn	
Spurry, clover	

## CABBAGE (Transplanted Only)

Table 2

<b>F7127 SE Herbicide Use Rate Table (Cabbage)</b>			
Fall or Spring Early Preplant, Preemergence, and Preplant Incorporated Applications			
<b>Broadcast Rate</b>	<b>Fl oz (lb ai) F7127 SE Herbicide per acre</b>		
	<b>Soil Texture</b>		
<b>% Organic Matter</b>	<b>Coarse</b>	<b>Medium</b>	<b>Fine</b>
<1.5%	2.9(0.08)- 3.8(0.10)	3.8(0.10)- 5.7(0.16)	3.8(0.10)- 7.6(0.21)
1.5-3.0 %	3.8(0.10)- 7.6(0.21)	7.6(0.21)- 11.4(0.31)	7.6(0.21)- 11.4(0.31)
>3.0 %	7.6(0.21)- 11.4(0.31)	7.6(0.21)- 15.2(0.41)	7.6(0.21)- 15.2(0.41)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories. Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.			

### Early Preplant (Fall Application or Spring Application)

F7127 SE Herbicide may be applied only in the fall or spring preceding the growing season to control weeds prior to or up to the planting or transplanting of cabbage. F7127 SE Herbicide may be applied in the spring from 60 days prior to planting up to planting time. F7127 SE Herbicide should be applied to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product into the soil. Do not mechanically incorporate in the fall or spring after application as this may destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils to prevent F7127 SE Herbicide runoff from rain or snow that may occur following application. F7127 SE Herbicide may be tankmixed with other burndown herbicides to control emerged weeds in the fall or spring or with residual soil herbicides that are labeled for fall use on cabbage. Use the full, listed rates of burndown herbicides in combination with F7127 SE Herbicide, or split applications as needed. Observe the most restrictive of all precautions, instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

### Preplant Incorporated (PPI)

F7127 SE Herbicide may be applied as a preplant incorporated treatment in the spring prior to transplanting of cabbage. Do not incorporate to depths greater than 2 inches. F7127 SE Herbicide can be tankmixed with other burndown or soil-applied herbicides labeled for use in cabbage. Use the full, listed rates of burndown herbicides or split applications as needed. Observe the most restrictive of all precautions, instructions and rotational cropping guidelines of each product's label when tank mixing including all references to potential carryover and crop injury warnings or restrictions.

### Transplant Cabbage

F7127 SE Herbicide may be applied pre-emergence as a broadcast or banded treatment to transplanted cabbage only. Make broadcast or banded treatment applications prior to transplanting. F7127 SE Herbicide may be applied as a banded treatment into the row middles within 72 hours after transplanting.

### Weeds Controlled

When Applied according to directions, F7127 SE Herbicide will provide control of:

Galinsoga, hairy	Waterhemp, common
Lambsquarters, common	Waterhemp, tall
Pigweed, redroot	

For information on other weeds not listed above, refer to WEEDS CONTROLLED section in this label.

### Precautions

These Crop Specific Use directions are based upon the interactive effects of F7127 SE Herbicide (sulfentrazone and carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE Herbicide Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been

evaluated under treatment with F7127 SE Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide under specific local conditions.

**Restrictions**

Do not apply more than 15.2 fluid ounces per acre of F7127 SE Herbicide per application or per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE Herbicide application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not incorporate to depths greater than 2 inches.

Pre-harvest Interval (PHI): 80 days

**CORN (Field Corn, Seed Corn, Popcorn)**

**Preplant Burndown, Early Preplant, and Preemergence Applications**

Apply F7127 SE HERBICIDE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to emergence of corn to control or suppress weeds using rates in table 3. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated with a preplant application of F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control. Apply F7127 SE HERBICIDE using the rates in Table 3 below.

For applications 14-21 or more days prior to planting, use the mid to high rate in the appropriate rate range for the soil and organic matter type listed in Table 3. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing F7127 SE HERBICIDE with other products be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

**Table 3**

<b>F7127 SE HERBICIDE Use Rate Table (Corn)</b>			
Preplant Burndown, Early Preplant, and Preemergence			
<b>Broadcast Rate</b>	<b>Fluid Ounces (lb ai) F7127 SE HERBICIDE per acre</b>		
	<b>Soil Texture</b>		
<b>% Organic Matter</b>	<b>Coarse</b>	<b>Medium</b>	<b>Fine</b>
<1.5	3.75(0.10)- 5.75(0.16)	3.75(0.10)- 5.75(0.16)	5.0(0.14)- 6.7(0.18)
1.5 – 3.0	3.75(0.10)- 5.75(0.16)	5.0(0.14)- 7.6(0.21)	5.75(0.16)- 8.6(0.23)
>3.0	5.0(0.14)- 7.6(0.21)	5.75(0.16)- 8.6(0.24)	7.6(0.21)- 10.2(0.28)
*Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

**Adjuvant Requirements**

For optimum control of emerged weeds a nonionic surfactant, crop oil concentrate, methylated seed oil, or equivalent adjuvant is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints/100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.5 to 2.0% v/v (1.5 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2.0 to 4.0% v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre may be used in addition to the selected NIS, COC, or MSO.

For all products used in tank mixes refer to the specific product labels for all restrictions on tank mixing and observe the most restrictive of all label precautions, instructions, and rotational cropping restrictions.

**Precautions**

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE HERBICIDE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions. Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor.

When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):

<b>BROADLEAVES</b>	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrisc</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Thistle, Russian	<i>Lactuca serriola</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
<b>SEDGES</b>	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

#### Restrictions

Do not apply F7127 SE HERBICIDE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snowmelt that may occur following application.

## DRY SHELLED BEANS AND PEAS

Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea moth bean, lentil, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lab lab bean; pea (*Pisum*) (includes dry field pea) and pigeon pea

#### Fall Applications

F7127 SE HERBICIDE may be applied in the fall as a preplant treatment to control or suppress weeds prior to planting the following spring. Apply F7127 SE HERBICIDE to the stubble or soil surface and allow moisture from rainfall or snow to move the product into the soil. Do not mechanically incorporate in the fall or spring as this can destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils or to existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snow melt that may occur following application. F7127 SE HERBICIDE may be tank mixed with other residual soil herbicides that are labeled for fall use on dry bean and dry peas. If weeds are emerged at the time of F7127 SE HERBICIDE application, use a burndown herbicide such as glyphosate or paraquat at the full-labeled rate in combination with F7127 SE HERBICIDE or split application as needed. Select the appropriate rate from the table below within the correct soil type and organic matter range. When applying F7127 SE HERBICIDE in the fall, use a mid to high rate within the rate range for the appropriate soil type and organic matter.

#### Preplant Burndown, Early Preplant, and Preemergence Applications

Apply F7127 SE HERBICIDE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or up to 3 days after planting dry shelled peas and beans to control or suppress weeds. Properly closed seed furrows are required when applying at planting time. When planting into soil treated preplant with F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing F7127 SE HERBICIDE with other products be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

**Table 4**

<b>F7127 SE HERBICIDE Use Rate Table (Dry Shelled Beans and Peas)</b>			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
<b>Broadcast Rate</b>	<b>Fl oz (lb ai) F7127 SE HERBICIDE per acre*</b>		
	<b>Soil Texture</b>		
<b>% Organic Matter</b>	<b>Coarse</b>	<b>Medium</b>	<b>Fine</b>
<1.5%	3.0(0.08) – 3.75(0.10)	3.75(0.1) – 5.75(0.16)	3.75(0.10) – 5.75(0.16)
1.5-3.0 %	3.75(0.10) – 5.75(0.16)	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 7.75(0.21)
>3.0 %	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 8.6(0.23)	6.7(0.18) – 10.2(0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

**Precautions**

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1.5%) and/or pH of 7.8 or higher, or on highly eroded soils (such as hilltops), or in areas of calcareous outcroppings. Reduce the use rates of F7127 SE HERBICIDE or do not apply it in those areas to avoid crop injury. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response. On coarse soils, best results are achieved with F7127 SE HERBICIDE when applications are made early preplant and greater than 14 days before planting.

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE HERBICIDE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions.

Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor. Optimum broad-spectrum control of annual and perennial weeds requires a tank-mix of with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat.

If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergent weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

**When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):**

Amaranth, Palmer	Pigweed, red root
Filaree, redstem	Pigweed, smooth
Kochia (ALS and Triazine Resistant)	Sida, prickly
Lambsquarters, common	Thistle, Russian
Morningglory, ivyleaf	Waterhemp, common
Morningglory, tall	Waterhemp, tall
Nightshade, Eastern black	

**Restrictions**

Do not apply F7127 SE HERBICIDE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or to existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snow melt that may occur following application.

## FALLOW SYSTEMS

F7127 SE HERBICIDE may be used in fallow cropping systems for weed control for soil moisture conservation using rates in Table 5. Fallow crop rotational restrictions when replanting following F7127 SE HERBICIDE applications.

Table 5

F7127 SE HERBICIDE Use Rate Table			
Fallow Applications			
Broadcast Rate	Fluid Ounces (lb ai) F7127 SE HERBICIDE per acre		
Soil Texture			
% Organic Matter	Coarse	Medium	Fine
<1.5	3.75(0.10)- 5.0(0.14)	3.75(0.10)- 5.75(0.16)	5.0(0.14)- 6.5(0.18)
1.5-3.0	3.75(0.10)- 5.75(0.16)	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.5(0.23)
>3	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.5(0.23)	6.5(0.18)- 10.2(0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

### Adjuvant Requirements

For optimum control of emerged weeds a nonionic surfactant, crop oil concentrate, methylated seed oil, or equivalent adjuvant is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints/100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.5 to 2.0% v/v (1.5 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2.0 to 4.0% v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre may be used in addition to the selected NIS, COC, or MSO. When an adjuvant is to be used with this product, FMC recommends use of a Chemical Producers and Distributors Association certified adjuvant.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing F7127 SE HERBICIDE with other products be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

For all products used in tank mixes refer to the specific product labels for all restrictions on tank mixing and observe the most restrictive of all label precautions, instructions, and rotational cropping restrictions.

### Precautions

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE HERBICIDE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions. **Thorough coverage is essential for control of small susceptible broadleaf weeds.** If thorough coverage is not achieved, postemergent weed control will be poor. If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergent weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

**When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):**

Kochia (ALS and Triazine Resistant)	Pigweed, redroot
Lambsquarters, common	Pigweed, smooth
Morningglory, ivyleaf	Thistle, Russian
Morningglory, tall	Waterhemp, common
Nightshade, Eastern Black	Waterhemp, tall

### Restrictions

Do not apply more than 10.2 fluid ounces per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snowmelt that may occur following application.

## FLAX

### APPLICATION TIMING - Fall Application, Early Preplant, and Preemergence Applications

#### FALL APPLICATION

F7127 SE HERBICIDE may be applied in the fall following crop harvest or in existing fallow fields to control or suppress weeds the following season. The F7127 SE HERBICIDE Rotational Crop Guidelines must be followed if crops are planted the next season. Apply F7127 SE HERBICIDE to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product. Do not mechanically incorporate in the fall or spring after application because this activity may destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils to prevent F7127 SE HERBICIDE runoff from rain or snow that may occur following application. F7127 SE HERBICIDE may be tank mixed with herbicides to control emerged weeds. Sequential applications of burndown herbicides may be needed depending on weed size. In situations where weed size may interfere with F7127 SE HERBICIDE reaching the soil surface, a separate burndown application prior to the application of F7127 SE HERBICIDE will be required. Use full, label-listed rates of burndown herbicides in combination with F7127 SE HERBICIDE, or sequential applications as needed. Higher aerial spray volumes are required when there is a dense weed population or canopy. Thorough coverage is essential for post-emergence control of small susceptible labeled broadleaf weeds in combination with glyphosate.

F7127 SE HERBICIDE can be tank mixed with other herbicides. Observe the most restrictive of all precautions, instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

#### SPRING APPLICATION - Early Preplant, and Preemergence Applications

Apply F7127 SE HERBICIDE alone or with other herbicides as a preemergence treatment prior to planting or up to 3 days after planting flax for preemergence control of susceptible broadleaf weeds using rates listed in the table below. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

When tank mixing F7127 SE HERBICIDE with other products be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 6

F7127 SE HERBICIDE Use Rate Table (Flax)			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fluid Ounces F7127 SE HERBICIDE per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	3.75 (0.10) – 5.0 (0.14)	3.75 (0.10) – 5.75 (0.16)	5.0 (0.14) – 6.5 (0.18)
1.5-3.0	3.75 (0.10) – 5.75 (0.16)	5.0 (0.14) – 7.75 (0.21)	5.75 (0.16) – 8.5 (0.23)
>3	5.0 (0.14) – 7.75 (0.21)	5.75 (0.16) – 8.5 (0.23)	7.75 (0.21) – 10.2 (0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

#### Precautions

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1.5%) and pH of 7.0 or higher, or on highly eroded soils, hill tops, or in areas of calcareous outcroppings. Reduce F7127 SE HERBICIDE use rates to 3.75 oz/A or do not apply in those areas to avoid crop injury. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE HERBICIDE Product Use Rates, Crop Rotational Intervals, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions.

### Weed Control

Thorough coverage is essential for control of small susceptible, emerged broadleaf weeds. If thorough coverage is not achieved, postemergence weed control will be poor. Optimum broad-spectrum post-emergent control of emerged weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate. Failure to achieve adequate burndown of existing vegetation prior to flax planting can result in poor crop growing conditions the remainder of the season. When tank-mixing F7127 SE Herbicide with other burndown herbicides for control of emerged weeds, it is recommended to use a full rate of the tank-mix herbicide. If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergence weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

When used as directed, F7127 SE HERBICIDE will provide preemergence control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergence weeds controlled):

Kochia (ALS and Triazine Resistant)	
Morningglory, ivyleaf*	
Morningglory, tall*	
Nightshade, Eastern black	

\* Partial or reduced control of the weeds listed above will occur under dry conditions, under heavy pest pressure or at low use rates under 5.75 oz.

### Restrictions

Do not apply F7127 SE HERBICIDE after crop emergence, or if the seedling is close to the soil surface as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre of F7127 SE HERBICIDE per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snowmelt that may occur following application.

## HORSERADISH

Table 7

<b>F7127 SE Herbicide Use Rate Table (Horseradish)</b>			
Fall or Spring Early Preplant, Preemergence, and Preplant Incorporated Applications			
<b>Broadcast Rate</b>	<b>Fl oz (lb ai) F7127 SE Herbicide per acre</b>		
	<b>Soil Texture</b>		
<b>% Organic Matter</b>	<b>Coarse</b>	<b>Medium</b>	<b>Fine</b>
<1.5%	2.9(0.08)- 5.7(0.16)	3.8(0.10)- 5.7(0.16)	3.8(0.10)- 5.7(0.16)
1.5-3.0 %	5.7(0.16)- 7.6(0.21)	7.6(0.21)- 10.2(0.28)	7.6(0.21)- 10.2(0.28)
>3.0 %	7.6(0.21)- 9.8(0.27)	7.6(0.21)- 10.2(0.28)	7.6(0.21)- 10.2(0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.			

F7127 SE Herbicide may be applied as an preplant premerge or preplant incorporated treatment by ground in a minimum of 15 gallons of finished spray.

### Early Preplant (Fall Application or Spring Application)

F7127 SE Herbicide may be applied in the fall or spring preceding the growing season to control or suppress weeds prior to or up to the planting of horseradish. F7127 SE Herbicide may be applied in the spring from 60 days prior to planting up to planting. Apply F7127 SE Herbicide to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product into the soil. Do not mechanically incorporate in the fall or spring after application as this may destroy the herbicide barrier and weed escapes may occur. Do not apply to frozen soils to prevent F7127 SE Herbicide runoff from rain or snow that may occur following application. F7127 SE Herbicide may be tankmixed with other burndown herbicides to control

emerged weeds in the fall or spring or with residual soil herbicides that are labeled for use on horseradish. Use full, listed rates of burndown herbicides in combination with F7127 SE Herbicide, or split applications as needed. Observe the most restrictive of all precautions, instructions, and rotational cropping guidelines of each product label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

**Preplant Incorporated (PPI)**

F7127 SE Herbicide may be applied as a preplant incorporated treatment in the spring prior to planting of horseradish. Do not incorporate to depths greater than 2 inches. F7127 SE Herbicide can be tankmixed with other burndown or soil-applied herbicides labeled for use on horseradish. Use the full, listed rates of burndown herbicides or split applications as needed. Observe the most restrictive of all precautions, instructions and rotational cropping guidelines of each product's label when tank mixing including all references to potential carryover and crop injury warnings or restrictions.

**Pre-Emergence (PRE)**

F7127 SE Herbicide may be applied pre-emergence as a broadcast or banded treatment on horseradish. Make broadcast applications prior to planting, or soon after planting but at least 5 days before crop emergence. F7127 SE Herbicide may be applied as a banded treatment into the row middles after crop emergence. Use the higher F7127 SE Herbicide rates on clay soils and/or soils with greater than 1% organic matter. F7127 SE Herbicide may be applied with other pesticides registered for use on horseradish.

**Weeds Controlled**

**When applied according to directions, F7127 SE Herbicide will provide control of:**

Lambsquarters, common	Pigweed, redroot
Morningglory, ivyleaf	Waterhemp, common
Nutsedge, yellow	Waterhemp, tall

For information on other weeds not listed above, refer to Weed Controlled section in this label.

**Precautions**

These Crop Specific Use directions are based upon the interactive effects of F7127 SE Herbicide (sulfentrazone and carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE Herbicide Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide under specific local conditions.

**Restrictions**

Do not apply more than 10.2 fluid ounces per acre of F7127 SE Herbicide per application or per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE Herbicide application.

Do not apply directly on the crop after the crop emerges or if the seedling sprouts are close to the soil surface.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not incorporate to depths greater than 2 inches.

**LIMA BEANS, SUCCULENT (Tennessee Only)**

**Preplant Burndown, Early Preplant, and Preemergence Applications .**

Apply F7127 SE HERBICIDE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting lima beans to control or suppress weeds. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

For applications 14-21 or more days prior to planting, use the mid to high rate in the appropriate rate range for the soil and organic matter type in Table 8. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing F7127 SE HERBICIDE with other products be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

**Table 8**

<b>F7127 SE HERBICIDE Use Rate Table (Lima Beans, Succulent)</b>			
Preplant Burndown, Early Preplant, and Preemergence			
<b>Broadcast Rate</b>	<b>Fl oz (lb ai) F7127 SE HERBICIDE per acre*</b>		
	<b>Soil Texture</b>		
<b>% Organic Matter</b>	<b>Coarse</b>	<b>Medium</b>	<b>Fine</b>
<1.5%	3.0 (0.08)- 5.0 (0.14)	3.8 (0.10)- 7.75 (0.21)	4.8 (0.13)-7.75 (0.21)
1.5 – 3.0 %	3.8 (0.10)- 5.75 (0.16)	5.0 (0.14)- 7.75 (0.21)	5.7 (0.16)-7.75 (0.21)
>3.0 %	5.0 (0.14)-	5.75 (0.16)-	6.7 (0.18)-7.75

	7.75 (0.21)	7.75 (0.21)	(0.21)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

**Precautions**

When applying F7127 SE HERBICIDE to coarse textured soils, growers must allow a minimum of 7-14 days from application to planting. Best results are achieved with F7127 SE HERBICIDE when applications are made early preplant and greater than 14 days before planting.

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1%) and pH of 7.8 or higher, or on highly eroded soils, or in areas of calcareous outcroppings. Reduce F7127 SE HERBICIDE use rates in those areas to avoid crop injury. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE HERBICIDE product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions.

Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor. Optimum broad-spectrum control of annual and perennial weeds requires a tank-mix of with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat.

**When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):**

Copperleaf, hophornbeam	Pigweed, redroot
Morningglory, entireleaf	Pigweed, smooth
Morningglory, ivyleaf	

**Restrictions**

Do not apply F7127 SE HERBICIDE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 7.75 fluid ounces per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or to existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snow melt that may occur following application.

Do not harvest forage or feed forage to livestock.

**PEANUTS**

**Southeastern United States Only (AL, AR, GA, LA, MS, NC, SC, TN, VA)**

**Preplant Burndown, Early Preplant, and Preemergence Applications.**

Apply F7127 SE HERBICIDE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or within 3 days after planting peanuts to control or suppress weeds using rates in Table 9. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing F7127 SE HERBICIDE with other products, be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

**Table 9**

<b>F7127 SE HERBICIDE Use Rate Table (Peanuts)</b>			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz (lb ai) F7127 SE HERBICIDE per acre*		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine

<1.5%	3.0(0.08) – 3.75(0.10)	3.75(0.10) – 5.75(0.16)	3.75(0.10) – 5.75(0.16)
1.5-3.0 %	3.75(0.10) – 5.75(0.16)	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 7.75(0.21)
>3.0 %	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 7.75(0.21)	6.5(0.18) – 10.2(0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

#### Precautions

When applying F7127 SE HERBICIDE with other registered herbicides, refer to specific label information on precautions, instructions, limitations, application methods and timings, and weeds controlled.

F7127 SE HERBICIDE is especially effective against a wide range of economic broadleaf weeds. The same processes that F7127 SE HERBICIDE affects in these weeds can, under certain conditions, be affected in peanuts. These conditions include high pH (7.5 and above), cool weather, prolonged and excessive moisture, seedling diseases, and any other condition, including poor agronomic practices, that are unfavorable to vigorous crop growth. Such effects in peanuts are often observed as stunting and discoloration. The duration of these effects are somewhat dependent on the duration of the adverse growing conditions. These effects lessen and generally diminish with a return to normal growing conditions. **Thorough coverage is essential for control of small susceptible broadleaf weeds.** If thorough coverage is not achieved, postemergent weed control will be poor

**When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):**

BROADLEAVES	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrisc</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Smartweed, PA (seedling)	<i>Polygonum pensylvanicum</i>
Thistle, Russian	<i>Lactuca serriola</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
SEDGES	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

#### Restrictions

Do not apply F7127 SE HERBICIDE Herbicide after crop emergence, at cracking, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre of F7127 SE HERBICIDE per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snowmelt that may occur following application. Do not apply after crop seed germination.

Do not feed treated peanut forage or peanut hay to livestock.

Do not irrigate with water having a pH higher than 9.

## POTATOES

Table 10

<b>F7127 SE Herbicide Use Rate Table (Potatoes)</b>			
Preemergence Application			
Broadcast Rate	Fl oz (lb ai) F7127 SE Herbicide per acre		
Soil Texture			
% Organic Matter	Coarse	Medium	Fine
<1.5	3.8(0.1)- 5.7(0.16)	3.8(0.10)- 5.7(0.16)	4.8(0.13)- 6.7(0.18)
1.5-3.0	3.8(0.10)- 5.7(0.16)	4.8(0.13)- 7.6(0.21)	5.7(0.16)- 7.6(0.21)
>3	5.7(0.16)- 7.6(0.21)	6.7(0.18)- 8.6(0.24)	7.6(0.21)- 10.2(0.28)

\*Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories  
Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.

### Ground and Aerial Applications

Apply F7127 SE Herbicide by aerial application as a preemergence treatment following planting and after dragoff, but prior to potato emergence. Optimum performance can be achieved if F7127 SE Herbicide is applied to the soil surface and either rainfall or overhead irrigation is used to activate the product. If no moisture is received within 7 days following application in areas without irrigation, a shallow incorporation (less than 2 inches) may be needed prior to weed and potato emergence to activate the product. Select the appropriate use rate based on soil texture and organic matter as shown in Table 10 above. For control of emerged weeds at the time of the F7127 SE Herbicide application, an appropriate burndown herbicide and adjuvants labeled for potatoes may be tankmixed with F7127 SE Herbicide to control these weeds. Do not apply F7127 SE Herbicide if the potatoes have emerged from the soil as undesirable crop response may occur. F7127 SE Herbicide may be tankmixed with other soil-applied herbicides labeled for use in potatoes to improve weed management and increase weed control spectrum.

Apply F7127 SE Herbicide in a minimum of 10 gallons of spray by ground application and 5 gallons of spray by air.

### Chemigation Applications

F7127 SE Herbicide may be applied to potatoes through sprinkler irrigation systems including center pivot, lateral move, end tow, solid set or hand move irrigation systems. Apply F7127 SE Herbicide prior to potato emergence using sufficient water (0.25 to 0.5 inch per acre) to provide thorough soil surface coverage, but to avoid runoff of irrigation water. F7127 SE Herbicide may be applied with other products labeled for chemigation use in potatoes.

It is important to note that irrigation with highly alkaline water (high pH) following a F7127 SE Herbicide soil application may significantly increase the amount of sulfentrazone available in soil solution. Irrigation with water having a pH greater than 7.5 could result in adverse crop response. This response will ultimately depend on initial F7127 SE Herbicide application rate, application timing, amount and pH of irrigation water; the sensitivity of the crop and the crop growth stage when irrigated. The risk of adverse crop response will lessen with advances in the crop growth stage.

### Weeds Controlled

When applied according to directions, F7127 SE Herbicide will provide control of:

Amaranth, Palmer	Nightshade, Eastern black
Filaree, redstem	Pigweed, redroot
Kochia (ALS and Triazine Resistant)	Pigweed, smooth
Lambsquarters, common	Thistle, Russian
Morningglory, ivyleaf	Waterhemp, common
Morningglory, tall	Waterhemp, tall

Also control all those weeds which are susceptible to carfentrazone application.

For information on other weeds not listed above, refer to Weed Controlled section in this label.

### Precautions

Potato varieties may vary in their response to herbicide applications. When using F7127 SE Herbicide on an untested variety, always determine the crop tolerance before planting. Some potato varieties, including Sangre, Shepody and Snowden, have shown sensitivity to F7127 SE Herbicide. Use extra caution when planting these varieties on marginal coarse soils.

These Crop Specific Use directions are based upon the interactive effects of F7127 SE Herbicide (sulfentrazone and carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE Herbicide Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label

pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide under specific local conditions.

### Restrictions

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply F7127 SE Herbicide after potato emergence from the soil as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE Herbicide application.

## SOYBEANS

### Preplant Burndown, Early Preplant, and Preemergence Applications .

Apply F7127 SE HERBICIDE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or within 3 days after planting soybeans to control or suppress weeds using rates in Table 11. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. F7127 SE HERBICIDE applied early pre-plant must be applied in combination with the appropriate burndown herbicide such as glyphosate, glufosinate, gramoxone, and/or 2,4-D to achieve acceptable control of existing weeds during application. When tank mixing F7127 SE HERBICIDE with other products, be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 11

<b>F7127 SE HERBICIDE Use Rate Table (Soybeans)</b>			
Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz (lb a.i.) F7127 SE HERBICIDE per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	5.75 (0.16)- 7.75 (0.21)	7.75 (0.21)-8.5 (0.23)	8.5 (0.23)
1.5-3	7.75 (0.21)- 8.5 (0.23)	8.5 (0.23)	8.5 (0.23)
>3	8.5 (0.23)	8.5 (0.23)	8.5 (0.23)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

### Precautions

When applying F7127 SE HERBICIDE with other registered herbicides, refer to specific label information on precautions, instructions, limitations, application methods and timings, and weeds controlled.

F7127 SE HERBICIDE is especially effective against a wide range of economic broadleaf weeds. The same processes that F7127 SE HERBICIDE affects in these weeds can, under certain conditions, be affected in soybeans. These conditions include high pH (7.5 and above), cool weather, prolonged and excessive moisture, seedling diseases, and any other condition, including poor agronomic practices, that are unfavorable to vigorous crop growth. Such effects in soybeans are often observed as stunting and discoloration. The duration of these effects are somewhat dependent on the duration of the adverse growing conditions. These effects lessen and generally diminish with a return to normal growing conditions. **Thorough coverage is essential for control of small susceptible broadleaf weeds.** If thorough coverage is not achieved, postemergent weed control will be poor.

**When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):**

BROADLEAVES	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrisc</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>

Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Smartweed, PA (seedling)	<i>Polygonum pennsylvanicum</i>
Thistle, Russian	<i>Lactuca scariola</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>
<b>SEDGES</b>	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

**Restrictions**

Do not apply F7127 SE HERBICIDE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 8.5 fluid ounces per acre of F7127 SE HERBICIDE per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snowmelt that may occur following application. Do not apply after crop seed germination. Do not feed soybean forage or hay to livestock.

**SUGARCANE**

Table 12

<b>F7127 SE Herbicide Use Rate Table (Sugarcane)</b>			
Planting Time and Lay-by Applications			
Broadcast Rate	Fl Oz (lb ai) per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	5.7 (0.16)- 7.6(0.21)	7.6(0.21)- 10.2(0.28)	10.2(0.28)
1.5-3	7.6(0.21)- 10.5(0.29)	10.2(0.28)- 12.8(0.35)	12.8(0.35)
>3	10.2(0.28)- 12.8(0.35)	12.8(0.35)- 15.2(0.41)	15.2(0.41)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.			

Apply F7127 SE Herbicide as a broadcast or banded preemergence soil applied treatment for the control of broadleaf weeds, grasses and sedges in sugarcane. Refer to the F7127 SE Herbicide Product Use Rate Section and Table 12 for specific use information.

**Planting Time Applications**

Apply F7127 SE Herbicide preemergence to newly planted or ratoon sugarcane. Use the higher rate on clay soils and/or soils with organic matter content higher than 2 percent. Apply either by air in a minimum of 5 gallons of spray per acre or by ground equipment in a minimum of 15 gallons of spray per acre. F7127 SE Herbicide may be applied with other herbicides registered for use in sugarcane.

**Lay-by Applications**

Apply F7127 SE Herbicide as a directed spray to sugarcane at lay-by timing. Use the higher rate on clay soils and/or soils with organic matter content higher than 2 percent. Apply as a directed spray with ground equipment in a minimum of 15 gallons of spray per acre. F7127 SE Herbicide may be applied with other herbicides registered for use in sugarcane.

**Weeds Controlled**

**When applied according to directions, F7127 SE Herbicide will provide control of:**

Morningglory, entireleaf	Morningglory, tall
Morningglory, ivyleaf	Pigweed, red root
Morningglory, red	Nutsedge, yellow

For information on other weeds not listed above, refer to WEEDS CONTROLLED section in this label.

### Precautions

These Crop Specific Use directions are based upon the interactive effects of F7127 SE Herbicide (sulfentrazone + carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE Herbicide Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide under specific local conditions.

### Restrictions

Pre-harvest Interval (PHI): Do not apply within 120 days of harvest.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not allow spray to contact crop leaves.

Do not apply more than 15.2 fluid ounces per acre of F7127 SE Herbicide per acre per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE Herbicide application.

## SUNFLOWERS

### Fall Application, Preplant Burndown, Early Preplant, and Preemergence Applications

#### Fall Application

F7127 SE HERBICIDE may be applied in the fall following crop harvest or in existing fallow fields to control or suppress weeds the following season. The F7127 SE HERBICIDE Rotational Crop Guidelines must be followed if crops are planted the next season. Apply F7127 SE HERBICIDE to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product. Do not mechanically incorporate in the fall or spring after application because this activity may destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils to prevent F7127 SE HERBICIDE runoff from rain or snow that may occur following application. F7127 SE HERBICIDE may be tank mixed with herbicides to control emerged weeds. Sequential applications may be needed depending on weed size. In situations where weed size may interfere with F7127 SE HERBICIDE reaching the soil surface, a separate burndown application prior to the application of F7127 SE HERBICIDE will be required. Use full, listed rates of burndown herbicides in combination with F7127 SE HERBICIDE, or sequential applications as needed. Higher aerial spray volumes are required when there is a dense weed population or canopy.

F7127 SE HERBICIDE can be tank mixed with other herbicides. Observe the most restrictive of all precautions, instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

#### Preplant Burndown, Early Preplant, and Preemergence Applications

Apply F7127 SE HERBICIDE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or up to 3 days after planting sunflowers to control or suppress weeds using rates in table 13. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with F7127 SE HERBICIDE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing F7127 SE HERBICIDE with other products be sure the F7127 SE HERBICIDE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 13

<b>F7127 SE HERBICIDE Use Rate Table (Sunflowers)</b>			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
<b>Broadcast Rate</b>	<b>Fl oz (lb ai) F7127 SE HERBICIDE per acre</b>		
	<b>Soil Texture</b>		
<b>% Organic Matter</b>	<b>Coarse</b>	<b>Medium</b>	<b>Fine</b>
<1.5	3.8(0.10)- 5.0(0.14)	3.8(0.10)- 5.75(0.16)	5.0(0.14)- 6.7(0.18)
1.5-3.0	3.8(0.10)- 5.75(0.16)	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.6(0.23)
>3	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.6(0.23)	7.75(0.23)- 10.2(0.28)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories  
Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.

**Precautions**

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1.5%) and pH of 7.8 or higher, or on highly eroded soils, hill tops, or in areas of calcareous outcroppings. Reduce F7127 SE HERBICIDE use rates or do not apply in those areas to avoid crop injury. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, F7127 SE HERBICIDE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions.

Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor. Optimum broad-spectrum control of annual and perennial weeds requires a tank-mix of with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat.

If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergent weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

**When used as directed, F7127 SE HERBICIDE will provide preemergent control of the following weeds (refer to POSTEMERGENT WEEDS CONTROLLED section for postemergent weeds controlled):**

Amaranth, Palmer	Pigweed, red root
Filaree, redstem	Pigweed, smooth
Kochia (ALS and Triazine Resistant)	Sida, prickly
Lambsquarters, common	Thistle, Russian
Morningglory, ivyleaf	Waterhemp, common
Morningglory, tall	Waterhemp, tall
Nightshade, Eastern black	

**Restrictions**

Do not apply F7127 SE HERBICIDE Herbicide after crop emergence, or if the seedling is close to the soil surface as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre of F7127 SE HERBICIDE per twelve-month period. The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

**Do not use on soils classified as sand, which have less than 1% organic matter.**

Do not apply to frozen soils or existing snow cover to prevent F7127 SE HERBICIDE runoff from rain or snowmelt that may occur following application.

**TOBACCO (Burley, Flue-Cured and Dark)**

Table 14

<b>F7127 SE Herbicide Use Rate Table (Tobacco)</b>			
Preemergence and Preplant Incorporated Applications			
Broadcast Rate	Fl oz (lb ai) F7127 SE Herbicide per acre		
Soil Texture			
% Organic Matter	Coarse	Medium	Fine
<1.5	5.7 (0.16)- 7.6(0.21)	7.6(0.21)- 10.2(0.28)	10.2(0.28)
1.5-3.0	7.6(0.21)- 10.2(0.28)	10.2(0.28)- 12.8(0.35)	12.8(0.35)
>3	10.2(0.28)- 12.8(0.35)	12.8(0.35)- 15.2(0.41)	15.2(0.41)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.			

F7127 SE Herbicide may be surface applied or preplant incorporated (to a depth no greater than 2 inches) from 14 days to 12 hours days prior to transplanting tobacco. Incorporating F7127 SE Herbicide deeper than 2 inches can result in inconsistent weed control. Broadcast apply the appropriate F7127 SE Herbicide rate from Table 14 above, in a minimum of 10 gallons per acre of water, to the soil prior to transplanting.

**Non-Bedded (Fields where raised beds are NOT formed prior to transplanting)**

Perform all accepted cultural practices for land preparation, fertilizer/fungicide incorporation, etc. prior to the application of F7127 SE Herbicide. Once the field has been prepared for planting, F7127 SE Herbicide may be surface applied or lightly preplant incorporated from 14 days to 12 hours prior to transplanting.

If F7127 SE Herbicide is surface applied and it is necessary to remove equipment tracks from the field after application but prior to transplanting, any light finishing equipment may be used providing the soil is not disturbed to a depth greater than 2 inches.

If timely cultivations are not performed following a pre-transplant surface application, reduced/unacceptable weed control may occur in the drill.

**Bedded (Fields where raised beds ARE formed PRIOR to transplanting)**

Apply F7127 SE Herbicide to formed beds as a surface application from 14 days to 12 hours prior to transplanting. If it is customary to drag/knock down beds prior to transplanting, this procedure must be performed prior to the F7127 SE Herbicide application.

When incorporating prior to bedding, F7127 SE Herbicide must be thoroughly and uniformly incorporated to a depth no greater than 2 inches to avoid concentrating F7127 SE Herbicide in the bed.

If initial transplanting fails to produce a uniform stand, tobacco may be replanted. DO NOT re-treat field with a second application of F7127 SE Herbicide, or any other herbicide containing sulfentrazone. DO NOT re-bed. Re-transplant into previously formed, treated beds.

For broad spectrum and optimum grass weed control a grass herbicide application will be required.

**Weeds Controlled**

When Applied according to directions, F7127 SE Herbicide will provide control of:
Filaree, redstem
Amaranthus, livid
Galinsoga, hairy
Lambsquarters, common
Morningglory, ivyleaf
Morningglory, tall
Pigweed, redroot
Pigweed, smooth
Sida, prickly
Signalgrass, broadleaf
Smartweed, Pennsylvania

For information on other weeds not listed above, refer to Weeds Controlled section in this label.

**Precautions**

Poor agronomic practices, unfavorable pH soils, diseases, cold weather, excessive moisture, drought or other conditions unfavorable to normal plant growth may adversely effect the growth of tobacco transplants. Weakened transplants may be more susceptible to herbicide response and diseases, particularly under poor drainage or compacted soil conditions or when the soil has been saturated for long periods of time. Contact your State Agricultural Extension Service Specialist for consultation as to the agronomic recommendations suited for your tobacco varieties and local conditions. Temporary stunting of tobacco may occur if transplants are set too shallowly, or if heavy rainfall occurs immediately following transplanting. Splashing of treated soil onto tobacco leaves may cause some localized and inconsequential necrosis. Use sound transplanting practices that insure treated soil will not wash or crust over tobacco plants.

These Crop Specific Use directions are based upon the interactive effects of F7127 SE HERBICIDE (sulfentrazone and carfentrazone-ethyl) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Application Instructions, Maximum Allowable F7127 SE Herbicide Use per Acre per 12 Month Period, Crop Rotational Intervals, Replanting Instructions, Weeds Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE HERBICIDE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE HERBICIDE under specific local conditions.

**Restrictions**

Do not use on Shade Grown Tobacco

Do not apply F7127 SE HERBICIDE to soils classified as sands containing less than 1% organic matter.

Do not use F7127 SE HERBICIDE in tobacco seeding beds or greenhouses.

Do not apply F7127 SE HERBICIDE post-transplant as unacceptable injury may occur.

Do not perform tillage practices that concentrate F7127 SE HERBICIDE into the bed or crop injury may occur.

Do not apply more than 15.2 fluid ounces (0.41 lbs active) per acre of F7127 SE HERBICIDE per acre per twelve-month period.

The twelve-month period is considered to begin upon the initial F7127 SE HERBICIDE application.

Do not incorporate greater than 2 inches deep.

## SOD PRODUCTION

F7127 SE Herbicide may be applied to established seeded, sodded or sprigged turfgrasses following the second mowing for the control of key grass, sedge and broadleaf weeds. Turf grasses should have developed a good root system, a uniform stand with healthy root systems to fill in the exposed edges prior to application. Sod injury could result from application of this product on sod that is not well established or has been weakened by stresses such as unfavorable weather conditions, diseases, chemical, recent harvesting or mechanical influences.

### Turf Grass Tolerance

When applied as directed, the following established turf grasses are tolerant to F7127 SE Herbicide herbicide at the listed use rates.

Table 15 Tolerant grasses

Grass Type	Maximum Use Rate For Single Application
<b>Cool Season Grasses **</b>	Fl oz (lb ai) F7127 SE Herbicide Per Acre
Bentgrass, creeping	5.1 (0.14)
Fescue, fine * (Festuca rubra) Fescue, tall * (Festuca arundinacea) Ryegrass, perennial (Lolium perenne) Bluegrass, Kentucky (Poa pratensis) Bluegrass, Rough (Poa trivialis)	5.1(0.14)-10.2(0.28)
<b>Warm Season Grasses **</b>	
Bahiagrass (Paspalum notatum) Buffalograss (Buchloe dactyloides) Carpetgrass (Axonopus affinis) Centipedegrass (Eremochloa ophiuoides) Kikuyugrass (Pennisetum clandestinum) Seashore Paspalum (Paspalum vaginatum) Zoysiagrass (Zoysia japonica) Bermudagrass (Cynadon dactylon) Bermudagrass Hybrids (Cyn Bluegrass, St. Augustinegrass (Stenotaphrum secundatum)	10.2(0.28)-15.2(0.41)

\* Applications of F7127 SE Herbicide to certain varieties of Chewings Fine Fescue or Tall Fescue may result in undesirable plant response.

\*\* It is important to note that not all varieties or cultivars have been evaluated under treatment with F7127 SE Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide under specific local conditions.

### Applications to Reseeded, Overseeded or Sprigged Areas

Reseeding, overseeding or sprigging may be done following F7127 SE Herbicide applications to turfgrasses. If reseeding, overseeding or sprigging is done within 1 month following a F7127 SE Herbicide treatment, the establishment of desirable grasses may be inhibited. Overseeding of bermudagrass with perennial ryegrass may be done two (2) to four (4) weeks following a F7127 SE Herbicide application provided slight grass plant response can be tolerated.

Optimum reseeding and overseeding results may be obtained with the use of mechanical or power seeding equipment, and where proper soil cultivation, irrigation and fertilization practices are followed.

### Adjuvant use

Good spray coverage is required for optimum control of weeds. Temporary discoloration of some sod species may result from use of surfactant. Use of surfactants is not recommended.

### Postemergence Control of Sedges

F7127 SE Herbicide may be applied at the rate of four (4) to twelve (12) fluid ounces per acre to established turf grasses for the control or suppression of sedges. Select the correct F7127 SE Herbicide use rate from Table 15

When applied as directed, F7127 SE Herbicide will provide control or suppression of the following sedges.

Table 16

Common Name	Scientific Name
Kyllinga, green	<i>Kyllinga brevifolia</i>
Kyllinga, false green	<i>Kyllinga gracillima</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, cylindrical	<i>Cyperus retrorsus</i>

Sedge, globe	<i>Cyperus globulosus</i>
Sedge, Surinam	<i>Cyperus surinamensis</i>
Sedge, Texas	<i>Cyperus polystachyos</i>

Purple nutsedge: For optimum control of purple nutsedge, split applications are listed below. Apply 4-8 ounces per acre as an initial application followed by a second application when evidence of actively growing purple nutsedge is visible. Do not exceed the maximum rate per acre based on the turf variety as listed in Table 15; tolerant grasses.

#### Split Application Rates for Optimum Purple Nutsedge Control

Grass Type	First Application (fl. oz (lb ai) per acre)	Second Application (fl. Oz (lb ai). per acre
Cool Season Grasses	2.5(0.07)-5.1(0.14)	2.5(0.07)-7.6(0.21)
Warm Season Grasses	5.1(0.14)-7.6(0.21)	5.1(0.14)-7.6(0.21)

Allow 35 days after first application for second application.

#### Postemergence Control of Grassy Weeds

F7127 SE Herbicide will control or suppress specific annual grasses (Table 17) when applied at a rate of 4 to 12 fl oz/acre . Apply the highest rate consistent with the rate needed for turfgrass tolerance in Table 15. Rates lower than 12 fl oz/acre will generally control grasses for at least 60 days. F7127 SE Herbicide works best if applied when the annual grasses are small (pre tiller stage) and actively growing.

**Table 17**

Common Name	Scientific Name
Goosegrass	<i>Eleusine indica</i>

#### Postemergence Control of Broadleaf Weeds

F7127 SE Herbicide herbicide will control or suppress the weeds listed in the broadleaf chart below when applied alone shortly after weeds have emerged. F7127 SE Herbicide may be applied at the rate of four (4) to twelve (12) fluid ounces per acre to established turf grasses for the control or suppression of broadleaf weeds. Select the correct F7127 SE Herbicide use rate from Table 15. For optimum results, make F7127 SE Herbicide applications shortly after weeds have emerged.

F7127 SE Herbicide may be tankmixed with other herbicides, insecticides and fungicides registered for use on turfgrasses. Read and follow the label directions and restrictions of the tank mix partner to determine turfgrass specie tolerance, use rates and application requirements. Follow all label restrictions, use directions and precautionary statements before use.

**When applied as directed, F7127 SE Herbicide will provide control or suppression of the following broadleaf weeds.**

Broadleaves	Scientific Names
Bittercress	<i>Cardamine spp.</i>
Black Medic	<i>Medicago lupulina</i>
Buttercup	<i>Ranunculus spp.</i>
Carolina geranium	<i>Geranium carolinianum</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Cinquefoil	<i>Potentilla spp.</i>
Clover	<i>Trifolium spp.</i>
Cudweed	<i>Gnaphalium spp.</i>
Dandelion	<i>Taraxacum officinale</i>
Dock, curly	<i>Rumex crispus</i>
Evening primrose	<i>Oenothera biennis</i>
Fiddleneck	<i>Amsinckia spp.</i>
Filaree	<i>Erodium spp.</i>
Garlic, wild	<i>Allium vineale</i>
Goldenrod	<i>Solidago spp.</i>
Ground ivy	<i>Glechema hederasea</i>
Henbit	<i>Lamium amplexicaule</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lawn burweed	<i>Soliva pterosperma</i>
Lespedeza, common	<i>Lespedeza striata</i>
Mallow, common	<i>Malva neglecta</i>
Onion, wild	<i>Allium canadense</i>
Parsley piert	<i>Alchemilla arvensis</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Pineapple weed	<i>Matricaria matricariodes</i>
Plantain, buckhorn	<i>Plantago lanceolata</i>
Puncture weed	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Redweed	<i>Melochia corchorifolia</i>
Rocket, London	<i>Sisymbrium irio</i>
Smartweed, PA	<i>Polygonum pensylvanicum</i>
Sorrel, red	<i>Rumex acetosella</i>

Speedwell	<i>Veronica spp.</i>
Spurge, annual	<i>Euphorbia spp.</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Star of Bethlehem	<i>Omithogalum umbellatum</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Violet, wild	<i>Viola pratincola</i>
Woodsorrel, creeping	<i>Oxalis corniculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>

#### Precautions

The use of additional surfactants may cause temporary undesirable effects to turfgrasses.

#### Restrictions

Sod production areas must be established three (3) months prior to the initial treatment of F7127 SE Herbicide.

Do not apply F7127 SE Herbicide to golf course greens or tees.

Do apply F7127 SE Herbicide to turf grasses not listed on this label.

Do not apply with surfactants without on-site evaluations for spray mixture compatibility and physical effects to turf grasses.

Do not graze or feed forage harvested from F7127 SE Herbicide treated areas.

Do not apply to landscape ornamental plants or ornamental beds.

Do not harvest sod within three (3) months of F7127 SE Herbicide application.

## PERMANENT CROPS:

### APPLES, CITRUS FRUIT, TREE NUTS, GRAPES and BERRIES

**Citrus Fruits (Crop Group 10):** Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these

**Grapes:** Wine, Raisin, Table and Juice, Amur river grape

**Berries (Crop Group 13-07):** aronia berry; bayberry; bearberry; bilberry; blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, ollieberry, Orgeon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these); blueberry, highbush; blueberry, lowbush; buffalo currant; buffaloberry; che; Chilean guava; chokecherry; cloudberry; highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); kiwifruit, fuzzy; kiwifruit, hardy; lingonberry; maypop; mountain pepper berries; mulberry; muntries; native currant; partridgeberry; phalsa; pincherry; rasperry, black and red; riberry; salal; schisandra berry; sea buckthorn; serviceberry; strawberry; wild raspberry; cultivars, varieties, and/or hybrids of these

**Tree Nuts (Crop Group 14):** Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (HazelNut), Hickory Nut, Macadamia Nut (Bush Nut), Pecan, Pistachio and Walnut (Black and English)

#### APPLICATION INFORMATION

Apply F7127 SE Herbicide as a uniform broadcast soil application to orchard, grove and vineyard floors and to berry beds and furrows or as a uniform band application directed to the base of the tree trunk and vines or beds in berries to provide preemergence control of weeds in table below.

For broadcast applications, make a single application of F7127 SE Herbicide at 7.7 to 15.2 fl oz per acre (0.21 to 0.41 lb ai/A). Do not apply more than 15.2 fluid ounces (0.41 lb ai) per acre per twelve-month period. The twelve-month period is considered to begin when the initial application of F7127 SE Herbicide is applied.

For improved weed management, F7127 SE Herbicide can be applied in a tank mixture with other preemergence and postemergence burndown herbicides. Refer to the tank mix partner's labels for additional restrictions, including minimum spray volumes and crops in which they are labeled. Burndown herbicides may include, but are not limited to, glyphosate, paraquat and glufosinate. Do not tank mix with Chateau® herbicides (flumioxazin) or with other products containing sulfentrazone.

When applied as a banded treatment (50% band or less), refer to formula in chart below for rate and volume. F7127 SE Herbicide may be applied twice per year. Do not apply more than 15.2 fl oz product (0.41 lb ai) per acre on a broadcast application basis per year. Allow a minimum of 60 days between applications.

For band treatments, apply the broadcast equivalent rate and volume per acre. To determine these:

Band Width Feet	X	Broadcast Rate Per Acre	=	Band Rate
Row Width Feet				
Band Width Feet	X	Broadcast Volume Per Acre	=	Band Volume

Use ground equipment only. Do not apply using an airblast sprayer or by air. Do not apply using a mechanically pressurized handgun.

Apply a minimum of 10 gallons of spray solution per acre to ensure uniform spray coverage. Nozzle selection must meet manufacturer's spray volume and pressure specifications for preemergence and postemergence herbicide applications. The spray solution should have a pH between 5.0 and 9.0.

Only apply F7127 SE Herbicide to crops that have been established for one full growing season and are in good health and vigor. Avoid contact of the spray solution on the green bark of trunks of 1-2 year old vines and trees by wrapping the trunk with a nonporous wrap, grow tubes, or wax containers which will keep the spray solution from coming in direct contact with the green tissue. Avoid direct or indirect spray contact with crop foliage and fruit.

Best results are obtained when the soil is moist at the time of application and allows for sufficient time for F7127 SE Herbicide to dry on the weed foliage prior to irrigation or rainfall and the application is followed by at least ½ inch of rainfall or sprinkler irrigation within two weeks after application. Time applications to take advantage of normal rainfall patterns and cool temperatures, especially where drip or micro sprinkler irrigation is used which may not uniformly incorporate the herbicide.

## WEED CONTROL INFORMATION

F7127 SE Herbicide provides burndown and is a selective soil-applied herbicide for the control of susceptible broadleaf, grass and sedge weeds found in Tables 18 and 19. Adequate moisture of ½ to 1 inch is required within 14 days after application for optimal control. If adequate rainfall is not received in a timely fashion, irrigate with a minimum of ½ inch of water. When activating moisture is delayed, a reduced level of weed control may occur. These escaped weeds can be removed using a burndown herbicide.

Tank mix F7127 SE Herbicide with a labeled postemergence burndown herbicide and use an appropriate adjuvant when weeds are present at the time of application. Refer to the tank mix partner's product label for the proper use rates by weed sizes. Use the most restrictive label limitations and precautions of the tank mix product(s).

Residual weed control may be reduced when F7127 SE Herbicide is applied where heavy crop trash such as leaves and branches and /or weed residues exists. It is best to rake or blow off the leaves and trash when they fall and prior to the F7127 SE Herbicide application.

Do not apply after petal fall (trees) or bud break (vines and berries) unless using a hooded or shielded sprayer to ensure that the spray solution will not come in contact with the crop or foliage.

### Permanent Crop Weed List

**Table 18**

Common Name	Scientific Name
Amaranth, livid	<i>Amaranthus lividus</i>
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus Powell II</i>
Amaranth, spiny	<i>Amaranthus spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Anoda, spurred	<i>Anoda cristata</i>
Barnyardgrass, common	<i>Echinochloa crus-galli</i>
Bedstraw, catchweed	<i>Galium aparine</i>
Bindweed, field	<i>Convolvulus arvensis</i>
Bluegrass, annual	<i>Poa annua</i>
Bromegrass species	<i>Bromus spp.</i>
Burclover, California	<i>Medicago polymorpha</i>
Carpetweed	<i>Mollugo verticillata</i>
Cheatgrass	<i>Bromus tectorum</i>
Cheeseweed species	<i>Malva spp.</i>

Chickweed, common	<i>Stellaria media</i>
Clover species	<i>Trifolium spp.</i>
Copperleaf, hophornbeam	<i>Acalypha ostryeafolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, Southern	<i>Digitaria ciliaris</i>
Croton, tropic	<i>Croton glandulosus</i>
Crownbeard, golden	<i>Verbesina encelioides</i>
Cupgrass, wooly	<i>Erichloa villosa</i>
Cyperus, hedgehog	<i>Cyperus compressus</i>
Daisy, American	<i>Eclipta alba</i>
Devilsclaw	<i>Proboscidea louisiana</i>
Dock, curly	<i>Rumex crispus</i>
Eclipta	<i>Eclipta prostrata</i>
Eveningprimrose, cutleaf	<i>Oenothera laciniata</i>
Fescue, Red	<i>Fetuca rubra</i>
Fiddleneck species	<i>Amsinckia spp.</i>
Filaree, broadleaf	<i>Erodium botrys</i>
Filaree, redstem	<i>Erodium cicutarium</i>
Filaree, whitestem	<i>Erodium moschatum</i>
Fleabane, hairy	<i>Conyza bonariensis</i>
Flixweed	<i>Descurainia sophia</i>
Foxtail, bristly	<i>Setari verticillata</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria glauca</i>
Galinsoga, hairy	<i>Galinsoga ciliata</i>
Goosegrass	<i>Eleusine indica</i>
Goosefoot, nettleleaf	<i>Chenopodium murale</i>
Groundcherry, clammy (seedling)	<i>Physalis heterophylla</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>
Groundsel, common	<i>Senecio vulgaris</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed (Marestail)	<i>Conyza canadensis</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass	<i>Sorghum halpense</i>
Junglerice	<i>Enchinochloa colona</i>
Knotweed, common	<i>Polygonum arenastrum</i>
Kochia (ALS and Triazine Resistant)	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, miners	<i>Montia perfoliata</i>
Lovegrass species	<i>Eragrostis spp.</i>
Mallow, common	<i>Malva neglecta wall r.</i>
Mallow, little	<i>Malva parviflora</i>
Mayweed, Chamomile	<i>Anthemis cotula l.</i>
Milkweed, honeyvine	<i>Ampelamus albidus</i>
Morningglory, entireleaf	<i>Ipomoea hederacea integruscula</i>
Morningglory, ivyleaf	<i>Ipomoea hederacea hederacea</i>
Morningglory, palmleaf	<i>Ipomoea wrightii</i>
Morningglory, purple	<i>Ipomoea turbinata</i>
Morningglory, red	<i>Ipomoea, coccinea L.</i>
Morningglory, scarlet	<i>Ipomoea coccinea</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomoea, purpurea</i>
Mullein, turkey	<i>Eremocarpus setigerus</i>
Mustard, Species	<i>Brassica spp.</i>
Mustard, tumble	<i>Sisymbrium altissimum</i>
Nettle, burning	<i>Urtica urens</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum ptycanthum</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Orchardgrass	<i>Dactylis glomerata</i>

Panicum, fall	<i>Panicum dichotomiflorum</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, Tumble	<i>Amaranthus albus</i>
Pineapple-weed	<i>Chamomilla suaveolens</i>
Plantain, blackseed	<i>Plantago rugelii decne</i>
Plantain, narrow-leaved	<i>Plantago lanceolata</i>
Poison Ivy	<i>Toxicodendron radicans</i>
Poorjoe	<i>Diodia teres</i>
Porophyllum	<i>Porophyllum rederule</i>
Poinsettia, wild	<i>Euphorbia heterophylla</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Redmaids	<i>Calandrinia ciliata</i>
Redweed	<i>Melochia corchorifolia</i>
Radish, Wild	<i>Raphanus raphanistrum</i>
Rocket, London	<i>Sisymbrium irio</i>
Sandbur	<i>Cenchrus spinifer</i>
Sedge, annual	<i>Carex spp.</i>
Senna, coffee	<i>Cassia occidentalis</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Sida, prickly	<i>Sida spinosa</i>
Sida, Southern	<i>Sida acuta</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, PA (seedling)	<i>Polygonum pennsylvanicum</i>
Smellmellon	<i>Cucumis melo</i>
Sowthistle species	<i>Sonchus spp.</i>
Sprangletop, red	<i>Leptochloa filiformis</i>
Spurge, spotted	<i>Chamaesyce maculate</i>
Starbur, bristly	<i>Acanthospermum hispidum</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Toadflax, yellow	<i>Linaria vulgaris</i>
Tassleflower, red	<i>Emilio sonchifolia</i>
Thistle, Russian	<i>Salsola kali</i>
Virginia Creeper	<i>Parthenocissus quiquefolia</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
Waterprimrose, winged	<i>Ludwigia decurrens</i>
Willowleaf, panicle-leaf	<i>Epilobium brachycarpum</i>
Witchgrass	<i>Panicum capillare</i>

#### ANNUAL AND PERENNIALSEDGE CONTROL INCLUDING NUTSEDGE

F7127 SE Herbicide applied at 15.2 ounces per acre may provide control or suppression of sedges whether applied preemergence or postemergence. Postemergence application to sedges allows F7127 SE Herbicide to be taken into the sedge through the foliage as well as soil uptake through the roots. Soil uptake is the major means of uptake by sedges. Good spray coverage is required for optimum control of sedges especially when applying postemergence to the sedges. Use a quality nonionic surfactant (NIS) at the rate of 0.25% v/v when applying postemergence.

When applied as directed, F7127 SE Herbicide will provide control or suppression of the following sedges.

Table 19

Common Name	Scientific Name
Kyllinga, green	<i>Kyllinga brevifolia</i>
Kullinga, false green	<i>Kyllinga gracillima</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, cylindrical	<i>Cyperus retrorsus</i>
Sedge, globe	<i>Cyperus globulosus</i>
Sedge, Surinam	<i>Cyperus surinamensis</i>
Sedge, Texas	<i>Cyperus polystachyos</i>

Optimum control of purple nutsedge may be obtained using split applications of F7127 SE Herbicide. Apply 5 – 7.7 ounces per acre followed by a second application to actively growing nutsedge. Do not exceed the maximum rate of 15.2 ounces (0.41 lb ai) per

season. F7127 SE Herbicide symptoms on nutsedge will be observed as reduced nutsedge stands, necrosis, chlorosis, and/or stunting. Optimum control may not be observed until the second year after the original treatment.

#### **REPLANTING IN NEW OR MATURE ORCHARDS AND VINEYARDS**

Delay replanting at least 30 days after F7127 SE Herbicide applications when replacing trees and vines in established orchards. Use untreated soil when replanting trees and vines.

#### **Precautions**

These Crop Specific Use directions are based upon the interactive effects of F7127 SE Herbicide Herbicide and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Product Application Instructions, F7127 SE Herbicide Herbicide Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE Herbicide Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide Herbicide under specific local conditions. FMC does not recommend tank mixing this product with other products containing sulfentrazone or other group 14 herbicides as crop injury may occur.

#### **Restrictions**

- Use ground equipment only. Do not apply F7127 SE Herbicide using airblast sprayers or by air. Do not apply using a mechanically pressurized handgun.
- Do not apply more than 15.2 oz product (0.41 lb ai total, 0.375 lb ai sulfentrazone and 0.124 lb ai carfentrazone-ethyl) per acre per 12-month season.
- Apply to crops that have been growing for at least one full year and are in good condition.
- Avoid direct or indirect spray contact to foliage and green bark (wrap trunk with non-porous wrap, grow tubes, or wax containers to keep spray solution off of green tissue).
- Do not apply to powdery soils or soils where wind may displace the soil, unless irrigation can be applied immediately after application.
- Follow the most restrictive label of tank mix partners including all references to potential carryover and crop injury warnings or restrictions.
- Pre-harvest Interval (PHI) for Apples only: 14 days
- Pre-harvest Interval (PHI) for Citrus Fruit, Tree Nuts, Grapes, and Berries : 3 days
- If two banded treatments are made in a growing season, allow a minimum of 60 days between applications ; however, do not exceed the seasonal maximum use rate.

## **LABEL TRACKING INFORMATION**

Label Code: 050515

FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia PA 19103

SPARTAN and FMC — Trademarks of FMC Corporation, Philadelphia, PA 19103 USA

©2015 FMC Corporation All Rights Reserved

# Supplemental Labeling

## F7127 SE Herbicide

EPA Reg. No.: 279-3337

**ACCEPTED**

05/18/2015

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under

EPA Reg. No. 279-3337

## DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

**This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for F7127 SE Herbicide before applying. Carefully follow all precautionary statements and application use directions.**

**This supplemental label expires June 1, 2018 and must not be used or distributed after this date.**

## APPLES

### APPLICATION INFORMATION

Apply F7127 SE Herbicide as a uniform broadcast soil application to orchard floors or as a uniform band application directed to the base of the tree trunk to provide preemergence control of weeds in table below.

For broadcast applications, make a single application of F7127 SE Herbicide at 7.7 to 15.2 fl oz per acre (0.21 to 0.41 lb ai/A). Do not apply more than 15.2 fluid ounces (0.41 lb ai) per acre per twelve-month period. The twelve-month period is considered to begin when the initial application of F7127 SE Herbicide is applied.

For improved weed management, F7127 SE Herbicide can be applied in a tank mixture with other preemergence and postemergence burndown herbicides. Refer to the tank mix partner's labels for additional restrictions, including minimum spray volumes and crops in which they are labeled. Burndown herbicides may include, but are not limited to, glyphosate, paraquat and glufosinate. Do not tank mix with Chateau® herbicides (flumioxazin) or with products containing sulfentrazone.

When applied as a banded treatment (50% band or less), refer to formula in chart below for rate and volume. F7127 SE Herbicide may be applied twice per year. Do not apply more than 15.2 fl oz product (0.41 lb ai) per acre on a broadcast application basis per year. Allow a minimum of 60 days between applications.

For band treatments, apply the broadcast equivalent rate and volume per acre. To determine these:

Band Width Feet	X	Broadcast Rate Per Acre	=	Band Rate
Row Width Feet				
Band Width Feet	X	Broadcast Volume Per Acre	=	Band Volume

Use ground equipment only. Do not apply using an airblast sprayer or by air. Do not apply using a mechanically pressurized handgun.

Apply a minimum of 10 gallons of spray solution per acre to ensure uniform spray coverage. Nozzle selection must meet manufacturer's spray volume and pressure specifications for preemergence and postemergence herbicide applications. The spray solution should have a pH between 5.0 and 9.0.

Only apply F7127 SE Herbicide to crops that have been established for one full growing season and are in good health and vigor. Avoid contact of the spray solution on the green bark of trunks of 1-2 year old vines and trees by wrapping the trunk with a nonporous wrap, grow tubes, or wax containers which will keep the spray solution from coming in direct contact with the green tissue. Avoid direct or indirect spray contact with crop foliage and fruit.

Best results are obtained when the soil is moist at the time of application and allows for sufficient time for F7127 SE Herbicide to dry on the weed foliage prior to irrigation or rainfall and the application is followed by at least ½ inch of rainfall or sprinkler irrigation within two weeks after application. Time applications to take advantage of normal rainfall patterns and cool temperatures, especially where drip or micro sprinkler irrigation is used which may not uniformly incorporate the herbicide.

## WEED CONTROL INFORMATION

F7127 SE Herbicide provides burndown and is a selective soil-applied herbicide for the control of susceptible broadleaf, grass and sedge weeds found in Tables 18 and 19. Adequate moisture of ½ to 1 inch is required within 14 days after application for optimal control. If adequate rainfall is not received in a timely fashion, irrigate with a minimum of ½ inch of water. When activating moisture is delayed, a reduced level of weed control may occur. These escaped weeds can be removed using a burndown herbicide.

Tank mix F7127 SE Herbicide with a labeled postemergence burndown herbicide and use an appropriate adjuvant when weeds are present at the time of application. Refer to the tank mix partner's product label for the proper use rates by weed sizes. Use the most restrictive label limitations and precautions of the tank mix product(s).

Residual weed control may be reduced when F7127 SE Herbicide is applied where heavy crop trash such as leaves and branches and /or weed residues exists. It is best to rake or blow off the leaves and trash when they fall and prior to the F7127 SE Herbicide application.

Do not apply after petal fall unless using a hooded or shielded sprayer to ensure that the spray solution will not come in contact with the crop or foliage.

### Permanent Crop Weed List

Common Name	Scientific Name
Amaranth, livid	<i>Amaranthus lividus</i>
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus Powell II</i>
Amaranth, spiny	<i>Amaranthus spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Anoda, spurred	<i>Anoda cristata</i>
Barnyardgrass, common	<i>Echinochloa crus-galli</i>
Bedstraw, catchweed	<i>Galium aparine</i>
Bindweed, field	<i>Convolvulus arvensis</i>
Bluegrass, annual	<i>Poa annua</i>
Bromegrass species	<i>Bromus spp.</i>
Burclover, California	<i>Medicago polymorpha</i>
Carpetweed	<i>Mollugo verticillata</i>
Cheatgrass	<i>Bromus tectorum</i>
Cheeseweed species	<i>Malva spp.</i>
Chickweed, common	<i>Stellaria media</i>
Clover species	<i>Trifolium spp.</i>
Copperleaf, hophornbeam	<i>Acalypha ostryeafolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, Southern	<i>Digitaria ciliaris</i>
Croton, tropic	<i>Croton glandulosus</i>
Crownbeard, golden	<i>Verbesina encelioides</i>
Cupgrass, wooly	<i>Erichloa villosa</i>
Cyperus, hedgehog	<i>Cyperus compressus</i>
Daisy, American	<i>Eclipta alba</i>
Devilsclaw	<i>Proboscidea louisiana</i>
Dock, curly	<i>Rumex crispus</i>
Eclipta	<i>Eclipta prostrata</i>
Eveningprimrose, cutleaf	<i>Oenothera laciniata</i>
Fescue, Red	<i>Fetuca rubra</i>
Fiddleneck species	<i>Amsinckia spp.</i>
Filaree, broadleaf	<i>Erodium botrys</i>
Filaree, redstem	<i>Erodium cicutarium</i>
Filaree, whitestem	<i>Erodium moschatum</i>

Fleabane, hairy	<i>Conyza bonariensis</i>
Flixweed	<i>Descurainia sophia</i>
Foxtail, bristly	<i>Setari verticillata</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria glauca</i>
Galinsoga, hairy	<i>Galinsoga ciliata</i>
Goosegrass	<i>Eleusine indica</i>
Goosefoot, nettleleaf	<i>Chenopodium murale</i>
Groundcherry, clammy (seedling)	<i>Physalis heterophylla</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>
Groundsel, common	<i>Senecio vulgaris</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed (Marestail)	<i>Conyza canadensis</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass	<i>Sorghum halpense</i>
Junglerice	<i>Enchinochloa colona</i>
Knotweed, common	<i>Polygonum arenastrum</i>
Kochia (ALS and Triazine Resistant)	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, miners	<i>Montia perfoliata</i>
Lovegrass species	<i>Eragrostis spp.</i>
Mallow, common	<i>Malva neglecta wall r.</i>
Mallow, little	<i>Malva parviflora</i>
Mayweed, Chamomile	<i>Anthemis cotula l.</i>
Milkweed, honeyvine	<i>Ampelamus albidus</i>
Morningglory, entireleaf	<i>Ipomoea hederacea integriuscula</i>
Morningglory, ivyleaf	<i>Ipomoea hederacea hederacea</i>
Morningglory, palmleaf	<i>Ipomoea wrightii</i>
Morningglory, purple	<i>Ipomoea turbinata</i>
Morningglory, red	<i>Ipomoea, coccinea L.</i>
Morningglory, scarlet	<i>Ipomoea coccinea</i>

Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomoea, purpurea</i>
Mullein, turkey	<i>Eremocarpus setigerus</i>
Mustard, Species	<i>Brassica spp.</i>
Mustard, tumble	<i>Sisymbrium altissimum</i>
Nettle, burning	<i>Urtica urens</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum ptycanthum</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Orchardgrass	<i>Dactylis glomerata</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, Tumble	<i>Amaranthus albus</i>
Pineapple-weed	<i>Chamomilla suaveolens</i>
Plantain, blackseed	<i>Plantago rugelii decne</i>
Plantain, narrow-leaved	<i>Plantago lanceolata</i>
Poison Ivy	<i>Toxicodendron radicans</i>
Poorjoe	<i>Diodia teres</i>
Porophyllum	<i>Porophyllum rederale</i>
Poinsettia, wild	<i>Euphorbia heterophylla</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Redmaids	<i>Calandrinia ciliata</i>
Redweed	<i>Melochia corchorifolia</i>

Radish, Wild	<i>Raphanus raphanistrum</i>
Rocket, London	<i>Sisymbrium irio</i>
Sandbur	<i>Cenchrus spinifer</i>
Sedge, annual	<i>Carex spp.</i>
Senna, coffee	<i>Cassia occidentalis</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Sida, prickly	<i>Sida spinosa</i>
Sida, Southern	<i>Sida acuta</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, PA (seedling)	<i>Polygonum pennsylvanicum</i>
Smellmellon	<i>Cucumis melo</i>
Sowthistle species	<i>Sonchus spp.</i>
Sprangletop, red	<i>Leptochloa filiformis</i>
Spurge, spotted	<i>Chamaesyce maculate</i>
Starbur, bristly	<i>Acanthospermum hispidum</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Toadflax, yellow	<i>Linaria vulgaris</i>
Tassleflower, red	<i>Emilio sonchifolia</i>
Thistle, Russian	<i>Salsola kali</i>
Virginia Creeper	<i>Parthenocissus quiquefolia</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
Waterprimrose, winged	<i>Ludwigia decurrens</i>
Willowleaf, panicle-leaf	<i>Epilobium brachycarpum</i>
Witchgrass	<i>Panicum capillare</i>

#### ANNUAL AND PERENNIALSEDGE CONTROL INCLUDING NUTSEDGE

F7127 SE Herbicide applied at 15.2 ounces per acre may provide control or suppression of sedges whether applied preemergence or postemergence. Postemergence application to sedges allows F7127 SE Herbicide to be taken into the sedge through the foliage as well as soil uptake through the roots. Soil uptake is the major means of uptake by sedges. Good spray coverage is required for optimum control of sedges especially when applying postemergence to the sedges. Use a quality nonionic surfactant (NIS) at the rate of 0.25% v/v when applying postemergence.

When applied as directed, F7127 SE Herbicide will provide control or suppression of the following sedges.

Common Name	Scientific Name
Kyllinga, green	<i>Kyllinga brevifolia</i>
Kullinga, false green	<i>Kyllinga gracillima</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, cylindrical	<i>Cyperus retrorsus</i>
Sedge, globe	<i>Cyperus globulosus</i>
Sedge, Surinam	<i>Cyperus surinamensis</i>
Sedge, Texas	<i>Cyperus polystachyos</i>

Optimum control of purple nutsedge may be obtained using split applications of F7127 SE Herbicide. Apply 5 – 7.7 ounces per acre followed by a second application to actively growing nutsedge. Do not exceed the maximum rate of 15.2 ounces (0.41 lb ai) per season. F7127 SE Herbicide symptoms on nutsedge will be observed as reduced nutsedge stands, necrosis, chlorosis, and/or stunting. Optimum control may not be observed until the second year after the original treatment.

#### REPLANTING IN NEW OR MATURE ORCHARDS AND VINEYARDS

Delay replanting at least 30 days after F7127 SE Herbicide applications when replacing trees and vines in established orchards. Use untreated soil when replanting trees and vines.

#### Precautions

These Crop Specific Use directions are based upon the interactive effects of F7127 SE Herbicide and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under Product Application Instructions, F7127 SE Herbicide Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with F7127 SE Herbicide. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on F7127 SE Herbicide under specific local conditions. FMC does not recommend tank mixing this product with other products containing sulfentrazone or other group 14 herbicides as crop injury may occur.

**Restrictions**

- Use ground equipment only. Do not apply F7127 SE Herbicide using airblast sprayers or by air. Do not apply using a mechanically pressurized handgun.
- Do not apply more than 15.2 oz product (0.41 lb ai total, 0.375 lb ai sulfentrazone and 0.124 lb ai carfentrazone-ethyl) per acre per 12-month season.
- Apply to crops that have been growing for at least one full year and are in good condition.
- Avoid direct or indirect spray contact to foliage and green bark (wrap trunk with non-porous wrap, grow tubes, or wax containers to keep spray solution off of green tissue).
- Do not apply to powdery soils or soils where wind may displace the soil, unless irrigation can be applied immediately after application.
- Follow the most restrictive label of tank mix partners including all references to potential carryover and crop injury warnings or restrictions.
- Pre-harvest Interval (PHI): 14 days
- If two banded treatments are made in a growing season, allow a minimum of 60 days between applications; however, do not exceed the seasonal maximum use rate.



FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia, PA 19103

Chateau® is a trademark of Valent USA Corporation

FMC is a trademark of FMC Corporation  
© 2015 FMC Corporation All Rights Reserved

050515